

REVIEW PAPER

A review on remediation technologies for dense metals polluted soil

Sadia Qayyum¹, Ibrar Khan³, Ke Meng¹, Yangguo Zhao¹, Changsheng Peng^{1,2,*}

¹The Key Lab of Marine Environmental Science and Ecology, Ministry of Education, Ocean University of China, Qingdao 266100, China

²State Key Laboratory of Environmental Criteria and Risk Assessment, Chinese Research Academy of Environmental Sciences, Beijing 100012, China


³UNESCO Chinese Center of Marine Biotechnology, Ocean University of China, Qingdao 266003, China



Highlights

- Due to industrialization and urbanization, environmental safety of soil has become a challenge.
- Biotechnological tools gradually became important technique for the last few decades for removal of metal ions pollution.
- Potassium phosphate is considered more effective in extracting arsenic among various potassium and sodium salts.
- The remediation mechanisms such as extracellular complexation, precipitation, oxidation-reduction reaction

Graphical Abstract



Article Info

Receive Date: 05 July 2019

Revise Date: 25 September 2019

Accept Date: 15 September 2019

Available online: 10 January 2020

Keywords:

Comparison
Contaminated soil
Environmental safety
Heavy metals
Remediation

Abstract

The Today, synthetic products such as industrial waste, pesticides, batteries, paints, and industrial or domestic sludge widely applied, as well as and manufacturing can adversely result in heavy metal contamination of urban and agricultural soils. Simultaneously, by growth of industrialization and urbanization, the ecological security of soil has become of great concerns. In view of examining the status of soil sullyng, the remediation advancements of soil debased by substantial metals were engaged in the current examination. To this point, physical remediation, concoction remediation and natural remediation were totally dedicated. To flexibly expected references to the current investigation, the instruments of remediation, qualities and downsides creating pattern were examined. It is suggested that for compelling and financial remediation of soil, a superior comprehension of remediation strategies and the different alternatives accessible at the various phases of remediation is exceptionally essential.



doi: 10.22034/CAJESTI.2020.01.03

* Corresponding author: pcs005@ouc.edu.cn (C. Peng)

E-ISSN: 2717-0519

P-ISSN: 2717-4034

1. Introduction

Soil is a natural and dynamic body formed by soil builder processes and factors, including minerals and organic materials that cover the earth's outer crust that plants can grow in it. The soil is considered essential natural components and significant material for individual endurance and improvement in the biological system. In any case, soil likewise turns into a path for contaminants entrance into the earth, disregarding defilement gets from air contamination, water contamination, or soil contamination itself (Fawzy et al., 2019; Khatun et al., 2019; Li et al., 2019; McBride and Zhou, 2019; Yu et al., 2019). Soil Contamination happens because of different horticultural and mechanical exercises, including petroleum product burning, agrarian utilization of manures and pesticides, mining waste, and landfill filtering (Almehdi et al., 2019; Azizollahi et al., 2019; Kumari and Dey, 2019). Potential unsafe contaminants have been collected in the higher soil during a great many years, beginning from the digging for haematite and far ahead for copper (Gong et al., 2019; Kumar et al., 2019; Parlayıcı and Pehlivan, 2019). Regarding their portability and bio harmfulness on living biological systems, expelling metals from soil is a fundamental undertaking. Soil tainting has become a significant issue with the economy and manufacturing improvement. Soil tainting by Heavy metal is more genuine than other soil defilements (Demarco et al., 2019; Gómez-Garrido et al., 2018; Guo et al., 2019; Jeelani et al., 2018; Liu et al., 2018).

By the aggregation of overwhelming metals and metalloids, soils sully may happen. Emanations from the quickly extending mechanical zones, removal of high metal squanders, mine tailings, leaded gas and paints, land use of composts, creature excrements, sewage muck, pesticides, wastewater water system, coal-burning buildups, spillage of petrochemicals, and environmental statement are of the chief reasons (Arreghini et al., 2018; Hossain et al., 2018; Khan et al., 2008; Lu et al., 2018; Tampouris et al., 2001; Zhang et al., 2018). Overwhelming metals are poisonous to all creatures if present in great obsessions. The convergences of substantial metals in soils can generally change, even in uncontaminated soils. Stamped contrasts in the stones' geochemical creation can bring about broad scopes of absolute centralizations of components in soils, even in soils not sullied. By the by, centralizations of substantial metals can influence harmfulness in soil creatures and vulnerable plants ward to the components influencing the bioavailability of the components (Ebadi and Hisoriev, 2017; Galal et al., 2019; Mai et al., 2019; Nan et al., 2019).

The traffic exhaust and fuel consumption, just as modern gases and residue, can prompt air contamination with many natural contaminants. The biological contaminants could then move to the dirt through different strategies, for example, dry testimony and precipitation, bringing about land tainting (Zhang, H. et al. 2017). In an exploration by (M.A. Hashim. et al., 2011), thirty-five methodologies for groundwater treatment have been assessed and ordered under three enormous classifications viz concoction, biochemical/natural/biosorption, and physical-substance treatment forms. The choice of an appropriate innovation for sully remediation at a specific site is one of the most testing activities because of very mind boggling soil science and spring attributes. No thumb-rule can be proposed for this issue. In the previous decade, iron-based advancements, microbial remediation, natural sulfate decrease and different adsorbents played flexible and proficient remediation jobs. Remembering the maintainability issues and ecological morals, the advancements including characteristic science, bioremediation and biosorption are prescribed to be embraced in proper cases.

2. Materials and Methods

2.1. Physical remediation

The physical remediation comprises of soil substitution strategy and warm desorption, too. The dirt substitution is to clean soil to reestablish or mostly reestablish the debased soil to weaken the contamination fixation, increment the natural dirt limit, and remediate the dirt (Abumaizar and Smith, 1999; Aresta et al., 2008; Kos and Leštan, 2003). The dirt substitution is grouped into three classes: soil substitution, soil spading, and new soil. Soil substitution expels the contaminated soil and brings in new soil, which is reasonable for a little debased zone. Furthermore, the supplanted soil ought to be dealt with potentially, or, more than likely, it will bring about the subsequent contamination. Soil spading is profoundly burrowing the defile soil, the toxin into the profound

locales, and accomplishing the point of weakening and normally debasing. New soil is dumping clean soil instead of the polluted soil, totally from base to surface or blending to diminish the poison fixation. The dirt supplanting is reasonable for soil with the little territory and contaminated harshly, which costs a ton (Abumaizar and Smith, 1999; Kos and Leštan, 2003). A few favorable circumstances, including necessary procedure, cell phones, and reuse capacity of the remediated soil, totally mean this innovation. An organization of mercury assortment and administration in USA has utilized this innovation for restoration and created business administration (Dandan et al., 2007).

2.2. Chemical remediation

2.2.1. Chemical leaching

Concoction draining is washing the sullied soil by utilizing new water, mixtures, and other liquids or gas (Tokunaga and Hakuta, 2002) to filter the contamination from the dirt. Sulfuric corrosive additionally accomplished excellent rate extraction. For arsenic expulsion from contaminated soil, a situation inviting and savvy extraction strategy has been contemplating (Alam et al., 2001; Lee et al., 2007). As an examined soil, a yellow-earthly colored woodland soil was sullied with arsenic (V) and utilized. Arsenic was capably extricated by phosphate arrangement of pH 6.0 at 300 mM phosphate focus and 40°C. The EDTA can shape constant complex with the most substantial metals in the wide pH valve among the extractants. (Ehsan et al., 2007). After one h of washing with 0.2 M citrus extract, expulsion efficiencies for fine silt were >95%. At the point when 0.2 M citrus extract was blende in with 0.1 M potassium phosphate, the as expulsion proficiency expanded to one hundred percent. Notably, the impact is practically unacceptable when solo extracting is utilized as a wide range of soil poisons. This lets us join or progressively use a wide range of extractants. As indicated by results, in the expulsion of overwhelming metals from the dirt examples, Na₂EDTA arrangements were profoundly more proficient than Na₂S₂O₅. Na₂EDTA especially showed a low effect on chromium evacuation and removed lead over zinc and cadmium. Evacuation of Cadmium and, significantly, zinc, by a 0.01 M Na₂EDTA arrangement, was expanded strikingly by incorporating 0.1 M Na₂S₂O₅. Consequently, a blend of the two reagents may give a monetarily ideal answer for firm debased soils (Ehsan et al., 2007).

Assessed the effectiveness of a coating procedure with cyclodextrin and EDTA for the concurrent activation of substantial metals and PCBs from a field-tainted soil. These examinations insisted that PCB blends and picked overpowering metals can be extracted capably from the soil with three dynamic washes with a comparative washing deferment having EDTA and cyclodextrin. Biodegradable, designed common chelate ethylenediaminedisuccinic destructive (EDDS) was used to wash soil polluted with 1350 mg/kg of Pb (Li et al., 2009). Additionally examined the proficiency of tea saponin on metal expulsion. The outcomes indicated that the evacuation of Pb, Cd, Zn, and Cu were 6.74, 42.38, 13.07, and 8.75%, separately when utilizing 7 wt % tea sapiens as the extracting. The tea sapiens can successfully evacuate corrosive solvent and reductive metals, which will extraordinarily lessen the ecological hazard.

2.2.2. Chemical fixation

Concoction obsession is to include mixtures or resources into the debased soil and to utilize them with substantial metals to frame impenetrable or barely portable, low poisonous issues, in this manner diminishing the movement of overwhelming metals to water, herb and other ecological media and to accomplish the remediation of soil (Abumaizar and Smith, 1999; Kos and Leštan, 2003). The outcomes demonstrated that Cd's centralization diminished 21.40, 27.63, 27.24, and 32.30% as contrasted and the regulator when the added substance sum was 20, 30, 50, and 40 g/kg, individually. There was likewise a report on the restoration of tainted soil by attapulgitic mud (Hong et al., 2002). Results exhibit that by including moderate attapulgitic dirt, Cd focus lessen 46% in soil, while the dirt profitability and quality were not influenced. Zhang et al. found that the compound obsession effectiveness of phosphate rock, furfural leftover, and endured coal on the polluted soil (Zhang et al., 2010). The consequences demonstrated that three molding operators could decrease the centralization of Cu, Zn, Pb, and Cd at specific grades. The substance obsession could remediate the dirt with low fixation pollutants; be that as it may,

the bioavailability of fixed substantial metals might be changed with the changing natural condition (Bolan et al., 2003). Also, molding specialists could change the dirt arrangement at certain amounts and impact the microorganisms in the soil.

2.2.3. *Electrokinetics remediation*

Electrokinetics remediation is another remediation innovation (Cabrera-Guzmán et al., 1990) in which voltage is applied at the different sides of soil, and afterward, the electronic ground angle is shaped. The poison was conveyed to two shafts cure room utilizing electro movement, electro-osmotic stream or electrophoresis, and later further treated (Virikutyte et al., 2002). It is appropriate for low penetrable soil, and has points of interest of effectively introduce and work, minimal effort (Cox et al., 1996; Virikutyte et al., 2002) and not pulverize the first nature condition (Cabrera-Guzmán et al., 1990; Hodson et al., 2000; Page and Page, 2002). so can reach the ecological restoration and secure the first ecotype (Cox et al., 1996). Be that as it may, the direct electrokinetics restoration can't control the pH estimation of soil framework well, and the dealing effectiveness was practically low. The principle enhanced techniques incorporate including support arrangement in cathode and anode to control pH esteem by utilizing particle trade layer to control pH esteem, to add complexant to improve movement, and so forth.

2.2.4. *Vitrify technology*

Vitrify innovation is to warm the dirt at a temperature of 1400~2000°C, wherein natural issues volatilize or disintegrate. The steam created and pyrolysis item was gathered by the off-gas dealing framework. In the wake of freezing, the liquefy structures rock profile glassy and make substantial metals lose movement. It was accounted for that the quality of the vitreous is multiple times higher than concrete. Petroleum product consuming or cathode straightforwardly warming provided required vitality for ex-situ remediation, and afterward, by circular segment, plasma and microwave vitality is moved. For in-situ restoration, the warmth can be through anodes embedded into the defiled soil. So, this innovation can expel the substantial metals, and productivity was high. Be that as it may, it is entangled and needs heaps of vitality in the softening, making it cost a great deal and restricted in the application (Zhang et al., 2010).

2.3. *Biological remediation*

The organic remediation incorporates Phytoremediation, bioremediation, and consolidating remediation.

2.3.1. *Phytoremediation*

From a global perspective, after the weather, the soil is considered the major component of the human environment. Soil not only is the major location for many of land creatures, especially human societies but also is a unique environment for the life of all species, especially plants. Unlike climate, soil contamination by chemical composition is not easily measurable and there is not a certain definition for pure soil, so we should study potential soil contamination issues in the context of predicting of hazards and potential damages in the soil function (Hong et al., 2002). Phytostabilization balances out substantial metals by plants through adsorption, rainfall and decrease of root. So that, root bioavailability lessen and forestall its relocating into the groundwater and evolved way of life (Hong et al., 2002).

Transgenic plants communicating marble developed progressively on a broad scope of convergences of monomethylmercuric chloride and phenylmercuric acetic acid derivation. Plants without the marble quality were carefully held or passed on at the equivalent organomercurial fixations. The study proposed that local macrophytes built to communicate merBpe might be utilized to debase methylmercury at dirtied locales and appropriate Hg (II) for later expulsion. In any case, this innovation is just reasonable for unpredictable contaminants, and the application is restricted (Hodson et al., 2000). Phytoextraction is absorbing the overwhelming metals utilizing lenient and gathering plants, and afterward moving, putting away. Considering the absorption depiction of diverse plants and selection, high take-up vegetation is the strategic of this advancement (Table 1).

Table 1. Some plant species that used in the phytoremediation process with the ability to absorb some heavy metals.

Plant Family	Plant Species	Metal
Araceae	<i>Pistia stratiotes</i>	Ag, Cd, Cr, Cu, Hg, Ni, Pb, and Zn
Asteraceae	<i>Helianthus indicus</i>	Pb
Fabaceae	<i>Sesbania drummondii</i>	Pb
Araceae	<i>Lemna gibba</i>	As
Solanaceae	<i>Solanum nigrum</i>	Cd
Brassicaceae	<i>Thlaspi caerulescens</i>	Cd

2.3.2 Biological remediation

However, the microorganisms can't corrupt and demolish the substantial metals, influencing the relocation and change over altering their physical and synthetic portrayals. The restoration systems incorporate extracellular complexation, rainfall, oxidation-decrease response, and intracellular amassing. An applied straightforward innovation for extricating valuable metals from second rate metals and mineral concentrates is Microbial filtering. Also, microbial draining can be possibly applied to remediation of mining destinations, detoxification of sewage slime, treatment of mineral modern waste items, and remediation of soils and residue sullied with substantial metals (Bosecker, 2001). Lambert et al. contemplated the impacts of muck on mycorrhizal (MR) take-up of P, Cu, and Zn, and affirm MR suppression of Cu and Zn take-up by P. Muck decreased take-up at 150 mg/kg P or higher in nonmycorrhizal (NMR) plants with little contrast in plant development among mucks (Lambert and Weidensaul, 1991). In any case, the natural restoration is vulnerable to influence by various types of elements, for example, temperatures, oxygen, dampness, pH esteem. Likewise, it was restricted in uses; for example, a few microorganisms can just debase exceptional contaminants, and organisms/zymin perhaps cause auxiliary contamination.

2.3.3. Animal remediation

Without a doubt, creature remediation is recognizing of some lower creatures adsorbing overwhelming metals, and afterward corrupting and relocating the substantial metals and along these lines expelling and restraining their harmfulness. The examinations indicated that the nightcrawler grass insulating blends upgraded plant Cu fixation, and the sum expanded by it was lower than that of the worm treatment yet more significant than that of straw mulching treatment (Lambert and Weidensaul, 1991). Zhang et al. examined the utilization of single extraction techniques to foresee the bioavailability of overwhelming metals in dirtied soils to rice. The outcomes demonstrated that substantial metals in soil could be estimated (Zhang et al., 2010). The collection sum expanded with the Pb focuses on developing.

3. Conclusion

The examination of remediation advancements is still at the personal and trial level. The advancement system of prospect remediation innovations examines green, natural amicable organic remediation, consolidating remediation, and in-situ remediation. To this point, speedy restoration and providing specialized support for farming soil tainting, mechanical ventures Brownfield, mining locales, etc. are to be applied. Soil contamination could influence crop profitability and human wellbeing. Exploring the sources, destiny, and event of soil contamination, and the dangers presented to human wellbeing has in this way been a significant territory of research.

Finally, plants and some algae for gathering environmental pollutions can also be important for human health in removing chemical methods that are very dangerous for environmental health and are costly and need more energy. This natural potential is unknown for many organisms and needs more and deep investigations to find better and optimal species. The local organisms show different treatments in different remediation methods in each specific ecological and environmental condition. It seems that many ways need to reach optimal ideas. Many plants and algae species in each country and the respective states can be used in refinery technology, so it can be

considered one of the research priorities in academic projects and dissertations. It can solve some of the environmental problems of the present age. In the end, It is important to note that the ecological and physiological characteristics of the given species have great importance. one of the processes that are used today in most countries is the need to provide a comprehensive atlas and detailed map of soil and fields pollution for the desired areas, so that, according to the type of pollution and its amount in each region, the optimal method will select for refining and sanitization.

Acknowledgments

The authors wish to recognize the necessary help from the administration, staff, and laborers of the Area Basalti srl whose full collaboration permitted the exploration program's improvement.

References

- Abumazar, R.J., Smith, E.H., 1999. Heavy metal contaminants removal by soil washing. *J. Hazard. Mater.*, **70**, 71–86. [https://doi.org/10.1016/S0304-3894\(99\)00149-1](https://doi.org/10.1016/S0304-3894(99)00149-1)
- Alam, M.G.M., Tokunaga, S., Maekawa, T., 2001. Extraction of arsenic in a synthetic arsenic-contaminated soil using phosphate. *Chemosphere*, **43**, 1035–1041. [https://doi.org/10.1016/S0045-6535\(00\)00205-8](https://doi.org/10.1016/S0045-6535(00)00205-8)
- Almehdi, A., El-Keblawy, A., Shehadi, I., El-Naggar, M., Saadoun, I., Mosa, K.A., Abhilash, P.C., 2019. Old leaves accumulate more heavy metals than other parts of the desert shrub *Calotropis procera* at a traffic-polluted site as assessed by two analytical techniques. *Int. J. Phytoremediation.*, **21**, 1254–1262. <https://doi.org/10.1080/15226514.2019.1619164>
- Aresta, M., Dibenedetto, A., Fragale, C., Giannoccaro, P., Pastore, C., Zammiello, D., Ferragina, C., 2008. Thermal desorption of polychlorobiphenyls from contaminated soils and their hydrodechlorination using Pd- and Rh-supported catalysts. *Chemosphere*, **70**, 1052–1058. <https://doi.org/10.1016/j.chemosphere.2007.07.074>
- Arreghini, S., de Cabo, L., Serafini, R.J.M., Fabrizio de Iorio, A., 2018. Shoot litter breakdown and zinc dynamics of an aquatic plant, *Schoenoplectus californicus*. *Int. J. Phytoremediation.*, **20**, 780–788. <https://doi.org/10.1080/15226514.2018.1425667>
- Azizollahi, Z., Ghaderian, S.M., Ghotbi-Ravandi, A.A., 2019. Cadmium accumulation and its effects on physiological and biochemical characters of summer savory (*Satureja hortensis* L.). *Int. J. Phytoremediation.*, **21**, 1241–1253. <https://doi.org/10.1080/15226514.2019.1619163>
- Bolan, N.S., Adriano, D.C., Duraisamy, P., Mani, A., 2003. Immobilization and phytoavailability of cadmium in variable charge soils. III. Effect of biosolid compost addition. *Plant Soil*, **256**, 231–241. <https://doi.org/10.1023/A:1026288021059>
- Bosecker, K., 2001. Microbial leaching in environmental clean-up programmes. *Hydrometallurgy*, **59**, 245–248. [https://doi.org/10.1016/S0304-386X\(00\)00163-8](https://doi.org/10.1016/S0304-386X(00)00163-8)
- Cabrera-Guzmán, D., Swartzbaugh, J.T., Weisman, A.W., 1990. The Use of Electrokinetics for Hazardous Waste Site Remediation. *J. Air Waste Manage. Assoc.*, **40**, 1670–1676. <https://doi.org/10.1080/10473289.1990.10466815>
- Cox, C.D., Shoesmith, M.A., Ghosh, M.M., 1996. Electrokinetic remediation of mercury-contaminated soils using iodine/iodide lixiviant. *Environ. Sci. Technol.*, **30**, 1933–1938. <https://doi.org/10.1021/es950633r>
- Dandan, W., Huixin, L., Feng, H., Xia, W., 2007. Role of earthworm-straw interactions on Phytoremediation of Cu contaminated soil by ryegrass. *Acta Ecol. Sin.*, **27**, 1292–1298. [https://doi.org/10.1016/S1872-2032\(07\)60030-4](https://doi.org/10.1016/S1872-2032(07)60030-4)
- Demarco, C.F., Afonso, T.F., Pieniz, S., Quadro, M.S., Camargo, F.A. de O., Andreatza, R., 2019. Phytoremediation of heavy metals and nutrients by the *Sagittaria montevidensis* into an anthropogenic contaminated site at Southern of Brazil. *Int. J. Phytoremediation.*, **21**, 1145–1152. <https://doi.org/10.1080/15226514.2019.1612843>
- Ebadi, A.G., Hisoriev, H., 2017. Metal pollution status of Tajan River – Northern Iran. *Toxicol. Environ. Chem.*, **99**, 1358–1367. <https://doi.org/10.1080/02772248.2017.1345191>

- Ehsan, S., Prasher, S.O., Marshall, W.D., 2007. Simultaneous mobilization of heavy metals and polychlorinated biphenyl (PCB) compounds from soil with cyclodextrin and EDTA in admixture. *Chemosphere*, **68**, 150–158. <https://doi.org/10.1016/j.chemosphere.2006.12.018>
- Fawzy, M., Nasr, M., Abdel-Rahman, A.M., Hosny, G., Odhafa, B.R., 2019. Techno-economic and environmental approaches of Cd²⁺ adsorption by olive leaves (*Olea europaea* L.) waste. *Int. J. Phytoremediation.*, **21**, 1205–1214. <https://doi.org/10.1080/15226514.2019.1612848>
- Galal, T.M., Shedeed, Z.A., Hassan, L.M., 2019. Hazards assessment of the intake of trace metals by common mallow (*Malva parviflora* K.) growing in polluted soils. *Int. J. Phytoremediation.*, **21**, 1397–1406. <https://doi.org/10.1080/15226514.2018.1524842>
- Gómez-Garrido, M., Mora Navarro, J., Murcia Navarro, F.J., Faz Cano, Á., 2018. The chelating effect of citric acid, oxalic acid, amino acids and *Pseudomonas fluorescens* bacteria on Phytoremediation of Cu, Zn, and Cr from soil using *Suaeda vera*. *Int. J. Phytoremediation.*, **20**, 1033–1042. <https://doi.org/10.1080/15226514.2018.1452189>
- Gong, Y., Chen, J., Pu, R., 2019. The enhanced removal and phytodegradation of sodium dodecyl sulfate (SDS) in wastewater using controllable water hyacinth. *Int. J. Phytoremediation.*, **21**, 1080–1089. <https://doi.org/10.1080/15226514.2019.1606779>
- Guo, Z., Gao, Y., Cao, X., Jiang, W., Liu, X., Liu, Q., Chen, Z., Zhou, W., Cui, J., Wang, Q., 2019. Phytoremediation of Cd and Pb interactive polluted soils by switchgrass (*Panicum virgatum* L.). *Int. J. Phytoremediation.*, **21**, 1486–1496. <https://doi.org/10.1080/15226514.2019.1644285>
- Hashim, M. A., Mukhopadhyay, S., Sahu, J. N., & Sengupta, B. 2011. Remediation technologies for heavy metal contaminated groundwater. *J. Environ. manage.*, **92**(10), <https://doi.org/2355-2388.10.1016/j.jenvman.2011.06.009>.
- Hodson, M.E., Valsami-Jones, É., Cotter-Howells, J.D., 2000. Bonemeal Additions as a Remediation Treatment for Metal Contaminated Soil. *Environ. Sci. Technol.*, **34**, 3501–3507. <https://doi.org/10.1021/es990972a>
- Hong, K.-J., Tokunaga, S., Kajiuchi, T., 2002. Evaluation of remediation process with plant-derived biosurfactant for recovery of heavy metals from contaminated soils. *Chemosphere*, **49**, 379–387. [https://doi.org/10.1016/S0045-6535\(02\)00321-1](https://doi.org/10.1016/S0045-6535(02)00321-1)
- Hossain, M.M., Khatun, M.A., Haque, M.N., Bari, M.A., Alam, M.F., Mandal, A., Kabir, A.H., 2018. Silicon alleviates arsenic-induced toxicity in wheat through vacuolar sequestration and ROS scavenging. *Int. J. Phytoremediation.*, **20**, 796–804. <https://doi.org/10.1080/15226514.2018.1425669>
- Jeelani, N., Yang, W., Qiao, Y., Li, J., An, S., Leng, X., 2018. Individual and combined effects of cadmium and polycyclic aromatic hydrocarbons on the phytoremediation potential of *Xanthium sibiricum* in co-contaminated soil. *Int. J. Phytoremediation.*, **20**, 773–779. <https://doi.org/10.1080/15226514.2018.1425666>
- Khan, S., Cao, Q., Zheng, Y.M., Huang, Y.Z., Zhu, Y.G., 2008. Health risks of heavy metals in contaminated soils and food crops irrigated with wastewater in Beijing, China. *Environ. Pollut.*, **152**, 686–692. <https://doi.org/10.1016/j.envpol.2007.06.056>
- Khatun, M.R., Mukta, R.H., Islam, M.A., Huda, A.K.M.N., 2019. Insight into citric acid-induced chromium detoxification in rice (*Oryza sativa* L.). *Int. J. Phytoremediation.*, **21**, 1234–1240. <https://doi.org/10.1080/15226514.2019.1619162>
- Kos, B., Leštan, D., 2003. Induced Phytoextraction/Soil Washing of Lead Using Biodegradable Chelate and Permeable Barriers. *Environ. Sci. Technol.*, **37**, 624–629. <https://doi.org/10.1021/es0200793>
- Kumar, V., AlMomin, S., Al-Shatti, A., Al-Aqeel, H., Al-Salameen, F., Shajan, A.B., Nair, S.M., 2019. Enhancement of heavy metal tolerance and accumulation efficiency by expressing *Arabidopsis* ATP sulfurylase gene in alfalfa. *Int. J. Phytoremediation.*, **21**, 1112–1121. <https://doi.org/10.1080/15226514.2019.1606784>

- Kumari, R., Dey, S., 2019. A breakthrough column study for removal of malachite green using coco-peat. *Int. J. Phytoremediation.*, **21**, 1263–1271. <https://doi.org/10.1080/15226514.2019.1633252>
- Lambert, D.H., Weidensaul, T.C., 1991. Element Uptake by Mycorrhizal Soybean from Sewage-Sludge-Treated Soil. *Soil Sci. Soc. Am. J.*, **55**, 393. <https://doi.org/10.2136/sssaj1991.03615995005500020017x>
- Lee, M., Paik, I.S., Do, W., Kim, I., Lee, Y., Lee, S., 2007. Soil washing of As-contaminated stream sediments in the vicinity of an abandoned mine in Korea. *Environ. Geochem. Health.*, **29**, 319–329. <https://doi.org/10.1007/s10653-007-9093-1>
- Li, J., Zhang, J., Larson, S.L., Ballard, J.H., Guo, K., Arslan, Z., Ma, Y., Waggoner, C.A., White, J.R., Han, F.X., 2019. Electrokinetic-enhanced Phytoremediation of uranium-contaminated soil using sunflower and Indian mustard. *Int. J. Phytoremediation.*, **21**, 1197–1204. <https://doi.org/10.1080/15226514.2019.1612847>
- Liu, X., Cao, L., Zhang, X., Chen, J., Huo, Z., Mao, Y., 2018. Influence of alkyl polyglucoside, citric acid, and nitrilotriacetic acid on Phytoremediation in pyrene-Pb co-contaminated soils. *Int. J. Phytoremediation.*, **20**, 1055–1061. <https://doi.org/10.1080/15226514.2018.1460305>
- Lu, Guangqiu, Wang, B., Zhang, C., Li, S., Wen, J., Lu, Guoli, Zhu, C., Zhou, Y., 2018. Heavy metals contamination and accumulation in submerged macrophytes in an urban river in China. *Int. J. Phytoremediation.*, **20**, 839–846. <https://doi.org/10.1080/15226514.2018.1438354>
- Mai, X., Luo, D., Wei, L., Liu, Y., Huang, X., Wu, Q., Yao, G., Liu, G., Liu, L., 2019. Evaluation method for the measuring comprehensive suitability of chelating agents: a study of the temporal dynamics of heavy metal activation. *Int. J. Phytoremediation.*, **21**, 1415–1422. <https://doi.org/10.1080/15226514.2019.1633262>
- McBride, M.B., Zhou, Y., 2019. Cadmium and zinc bioaccumulation by *Phytolacca americana* from hydroponic media and contaminated soils. *Int. J. Phytoremediation.*, **21**, 1215–1224. <https://doi.org/10.1080/15226514.2019.1612849>
- Nan, G., Guo, L., Gao, Y., Meng, X., Zhang, L., Song, N., Yang, G., 2019. Speciation analysis and dynamic absorption characteristics of heavy metals and deleterious element during growing period of Chinese peony. *Int. J. Phytoremediation.*, **21**, 1407–1414. <https://doi.org/10.1080/15226514.2019.1633261>
- Nicholson, F.A., Smith, S.R., Alloway, B.J., Carlton-Smith, C., Chambers, B.J., 2003. Quantifying heavy metal inputs to agricultural soils in England and Wales. *Water Environ. J.*, **20**, 87–95. [https://doi.org/10.1016/S0048-9697\(03\)00139-6](https://doi.org/10.1016/S0048-9697(03)00139-6)
- Page, M.M., Page, C.L., 2002. Electroremediation of Contaminated Soils. *J. Environ. Eng.*, **128**, 208–219. [https://doi.org/10.1061/\(ASCE\)0733-9372\(2002\)128:3\(208\)](https://doi.org/10.1061/(ASCE)0733-9372(2002)128:3(208))
- Parlayıcı, Ş., Pehlivan, E., 2019. Fast decolorization of cationic dyes by nano-scale zero valent iron immobilized in sycamore tree seed pod fibers: kinetics and modelling study. *Int. J. Phytoremediation.*, **21**, 1130–1144. <https://doi.org/10.1080/15226514.2019.1606786>
- Tampouris, S., Papassiopi, N., Paspaliaris, I., 2001. Removal of contaminant metals from fine grained soils, using agglomeration, chloride solutions and pile leaching techniques. *J. Hazard. Mater.*, **84**, 297–319. [https://doi.org/10.1016/S0304-3894\(01\)00233-3](https://doi.org/10.1016/S0304-3894(01)00233-3)
- Tokunaga, S., Hakuta, T., 2002. Acid washing and stabilization of an artificial arsenic-contaminated soil. *Chemosphere*, **46**, 31–38. [https://doi.org/10.1016/S0045-6535\(01\)00094-7](https://doi.org/10.1016/S0045-6535(01)00094-7)
- Virkutyte, J., Sillanpää, M., Latostenmaa, P., 2002. Electrokinetic soil remediation — critical overview. *Sci. Total Environ.*, **289**, 97–121. [https://doi.org/10.1016/S0048-9697\(01\)01027-0](https://doi.org/10.1016/S0048-9697(01)01027-0)
- Yu, F., Li, Y., Li, F., Zhou, Z., Chen, C., Liang, X., Li, C., Liu, K., 2019. Nitrogen fertilizers promote plant growth and assist in manganese (Mn) accumulation by *Polygonum pubescens* Blume cultured in Mn tailings soil. *Int. J. Phytoremediation.*, **21**, 1225–1233. <https://doi.org/10.1080/15226514.2019.1619161>

Zhang, J., Yang, S., Yang, H., Huang, Y., Zheng, L., Yuan, J., Zhou, S., 2018. Comparative study on effects of four energy plants growth on chemical fractions of heavy metals and activity of soil enzymes in copper mine tailings. *Int. J. Phytoremediation.*, **20**, 616–623. <https://doi.org/10.1080/15226514.2017.1413328>

Zhang, M.-K., Liu, Z.-Y., Wang, H., 2010. Use of Single Extraction Methods to Predict Bioavailability of Heavy Metals in Polluted Soils to Rice. *Commun. Soil Sci. Plant Anal.*, **41**, 820–831. <https://doi.org/10.1080/00103621003592341>

Zhang, H., Ma, D., Qiu, R., Tang, Y., Du, C. 2017. Non-thermal plasma technology for organic contaminated soil remediation: A review. *Chem. Eng. J.*, **313**, 157-170. <https://doi.org/10.1016/j.cej.2016.12.067>

Copyright © 2020 by CAS Press (Central Asian Scientific Press) + is an open access article distributed under the Creative Commons Attribution License (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

How to cite this paper:

Qayyum, S., Khan, I, Meng, K., Zhao, Y., Peng, C., 2020. A review on remediation technologies for heavy metals contaminated soil. *Cent. Asian J. Environ. Sci. Technol. Innov.* **1**(1), 21-29.